

# Stereoselective photodimerisation of chalcones in the molten state

Fumio Toda,<sup>\*,a</sup> Koichi Tanaka<sup>a</sup> and Masako Kato,<sup>\*,b</sup>

<sup>a</sup> Department of Applied Chemistry, Faculty of Engineering, Ehime University, Matsuyama, Ehime 790, Japan

<sup>b</sup> Department of Chemistry, Faculty of Science, Nara Women's University, Nara, Nara 630, Japan

Photodimerisations of chalcone and its derivatives in the molten state proceed efficiently and stereoselectively to give *rac-anti*-head-to-head dimers in all cases tested.


## Introduction

Although photodimerisations of chalcone and its derivatives in solution have long been studied, most gave unsatisfactory results. For example, irradiation of chalcone **1a**<sup>1</sup> and 4'-methylchalcone **1b**<sup>2</sup> in solution gave the corresponding *anti*-head-to-head dimers **5a** and **5b**, respectively, in very poor yield.<sup>3</sup> Photodimerisation of chalcones **1** in the solid state has also long been studied in order to correlate crystal-packing geometry with photochemical behaviour.<sup>4</sup> In spite of many efforts at photodimerisation by several research groups, all such attempts have failed except for one uncertain dimerisation experiment on 4-methoxychalcone **1d**<sup>5</sup> in the solid state which gave a mixture containing *syn*-head-to-head dimer **2d**.<sup>3</sup> X-Ray analysis of compound **1d** as a crystal showed that the shortest contact of C=C groups of two neighbouring **1d** molecules is 4.2 Å. In all other photochemically inert chalcones, the shortest contacts are longer than 4.2 Å. Finally, the rule 'the photodimerisation of chalcone occurs when the distance between C=C groups is shorter than 4.2 Å', the so-called Schmidt's rule, has appeared.<sup>6</sup> However, in an inclusion crystal of chalcone **1a** with 1,1,6,6-tetraphenylhexa-2,4-diyne-1,6-diol host compound **3**,<sup>7</sup> **1a** molecules aggregate in close positions and give product **4a** in 90% yield by photoirradiation in the solid state.<sup>7,8</sup>


## Results and discussion


We found that photodimerisations of chalcone and its derivatives proceed efficiently and stereoselectively in the molten state and give the corresponding *rac-anti*-head-to-head dimers **5** in relatively good yields. Irradiation of one polymorphic crystal of chalcone **1a** (form II, mp 56 °C)<sup>9</sup> (1 g) at 60 °C for 24 h by a 400 W high-pressure Hg lamp gave, after recrystallisation from MeOH, compound **5a** as prisms (0.31 g, 31%; mp 123–125 °C). Similar irradiation of another polymorphic crystal form of chalcone **1a** (form I, mp 59 °C)<sup>10</sup> at 60 °C also gave compound **5a** in 28% yield. This is a very efficient photodimerisation method, because neither form I nor form II of chalcone **1a** gives any dimer by photoirradiation in the solid state.<sup>9,10</sup> The structure of product **5a** was elucidated from its <sup>1</sup>H NMR spectrum which shows highly shielded CH proton signals at δ 3.97 and 4.61. The structure was finally determined by X-ray analysis (Fig. 1). Although X-ray data for compound **5a** have already been reported,<sup>11</sup> we used our own data, because of its improved quality. Dimer **5a** has been prepared so far in 9% yield as a crystal (mp 124 °C) by photoirradiation of chalcone **1a** in EtOH for 147 h, and the structure has been estimated by a chemical method.<sup>2</sup> Similar irradiation of compound **1b** (1 g, mp 74–75 °C) at 80 °C for 24 h gave, after recrystallisation from MeOH, dimer **5b** as prisms (0.27 g, 27%; mp 112–114 °C). By comparison of its <sup>1</sup>H NMR spectrum with that of dimer **5a**, the structure of compound **5b** was determined. By prolonged irradiation of compound **1b** in EtOH for 16 days, a mixture of its two kinds of dimers, mps 114 and 218 °C, has been obtained in 45% yield.<sup>9</sup> The former product would be identical with dimer **5b**. Similarly, irradiation of chalcones **1c** (mp 96–98 °C),<sup>11</sup> **1d** (mp 74 °C)<sup>5</sup> and **1e** (mp 127 °C)<sup>12</sup> in their molten state gave the corresponding *rac-anti*-head-to-head dimers **5c** (25% yield; mp 128–129 °C), **5d** (20% yield; mp 90–92 °C) and **5e** (32% yield; mp 137–139 °C), respectively, in the yields indicated. Structures of compounds **5c–e** were elucidated by comparison of their <sup>1</sup>H NMR spectra with that of the parent dimer **5a**.


All photodimerisations of chalcones **1a–e** in the molten state gave *rac-anti*-head-to-head dimers **5a–e**. This result suggests that two molecules of chalcone **1** aggregate in the liquid state so as to give the dimer **5** by photoreaction as shown in structure **6**. Since irradiation of substrates **1a** and **1b** in solution also gives dimers **5a** and **5b**, respectively, in low yields,<sup>2</sup> molecules of chalcone **1** would also be relatively easy to aggregate as species **6** even in solution. The aggregation of dimers **5a–e** as species **6** would be energetically the most favourable. Since photodimerisation of chalcones **1** is more efficient in the molten state than in solution, molecules of substrate **1** would more easily aggregate as structures **6** in the molten state. In solution, solvent molecules probably interrupt the molecular aggregation of chalcones **1**.





a; Ar = Ar' = Ph

b; Ar = Ph, Ar' = 

c; Ar = Ph, Ar' = 


d; Ar = , Ar' = Ph

e; Ar = Ar' = 




**Fig. 1** Molecular structure of dimer **5a** with crystallographic numbering scheme. Selected bond lengths (Å) and angles (°): O(1)–C(5) = 1.219(3), O(2)–C(12) = 1.216(3), C(1)–C(2) = 1.568(3), C(1)–C(4) = 1.541(3), C(1)–C(5) = 1.509(3), C(2)–C(3) = 1.567(3), C(2)–C(19) = 1.507(3), C(3)–C(4) = 1.563(3), C(3)–C(25) = 1.502(3), C(4)–C(12) = 1.509(3), C(5)–C(6) = 1.488(3), C(12)–C(13) = 1.496(3); C(2)–C(1)–C(4) = 90.0(2), C(2)–C(1)–C(5) = 116.1(2), C(4)–C(1)–C(5) = 116.2(2), C(1)–C(2)–C(3) = 89.4(2), C(1)–C(2)–C(19) = 120.1(2), C(3)–C(2)–C(19) = 114.7(2), C(2)–C(3)–C(4) = 89.2(2), C(2)–C(3)–C(25) = 116.8(2).


All of the molecules of chalcones **1a–e** aggregate similarly in their molten state as discussed above. However, molecules of chalcones **1** in crystals do not aggregate in a definite manner. The crystal structure of 4-methoxychalcone **1d** (mp 74 °C)<sup>5</sup> has been shown by X-ray analysis to have a molecular aggregation with 4.2 Å shortest C=C contact, which should give *syn*-head-to-head dimer **2d** by photodimerisation.<sup>3</sup> On the other hand, molecules of 4'-chlorochalcone **1c** were found to be arranged in the crystal in a different manner from that of analogue **1d**. Irradiation of powdered compound **1c**<sup>12</sup> (1 g) in a stirred water suspension containing a small amount of hexadecyl(trimethyl)ammonium bromide as a surfactant for 12 h by 100 W high-pressure Hg-lamp gave *syn*-head-to-tail dimer **4c** as prisms, after recrystallisation from AcOEt (0.31 g, 31%; mp 255–257 °C). An X-ray analysis showed that two **1c** molecules are aggregated at 4.238 Å shortest C=C contact in the positions which give the **4c** dimer by photoreaction (Fig. 2). However, 4'-methylchalcone **1b** molecules are arranged in the same way as are the **1c** substrate molecules, although C=C contact is too long, 4.5 Å, to dimerise in the solid state (Fig. 3). Chiral order of 4,4'-dimethylchalcone molecules **1e**<sup>13</sup> in their crystal state has been elucidated by X-ray analysis.<sup>13</sup> Enantioselective bromine addition to the chiral crystal of compound **1e** which gives the optically active bromine addition product in low optical yield has also been reported.<sup>14</sup> Although irradiation of the chiral crystal of compound **1e** was carried out in the solid state in the expectation of the formation of optically active photodimer, substrate **1e** was photochemically inert. This is very reasonable, because the C=C contact in the compound **1e** crystal is too long (5.9 Å) to react.<sup>13</sup> The chiral arrangement of compound **1e** molecules in the crystal was easily detected by measurement of CD spectra (Fig. 4) in Nujol mull.<sup>15</sup> In Fig. 4, a nice mirror-image relationship of two enantiomeric chiral crystals of **1e** is shown.



**Fig. 2** Molecular structure of compound **1c** with crystallographic numbering scheme. Selected bond lengths (Å) and angles (°): C(1)–C(7) = 1.736(3), O(1)–C(1) = 1.227(3), C(1)–C(2) = 1.475(4), C(1)–C(4) = 1.492(4), C(2)–C(3) = 1.323(4), C(3)–C(10) = 1.455(4); O(1)–C(1)–C(2) = 121.4(2), O(1)–C(1)–C(4) = 119.5(2), C(2)–C(1)–C(4) = 119.1(2), C(1)–C(2)–C(3) = 121.7(3), C(2)–C(3)–C(10) = 128.0(3). Non-bonded contacts (Å): C(2)⋯C(2') = 4.002(4), C(3)⋯C(3') = 4.840(4) and C(2)⋯C(3') = 4.239(4) (*l*' = 1 – *x*, –*y*, 1 – *z*).




**Fig. 3** Molecular structure of compound **1b** with crystallographic numbering scheme. Selected bond lengths (Å) and angles (°): O(1)–C(1) = 1.229(2), C(1)–C(2) = 1.478(2), C(1)–C(4) = 1.488(2), C(2)–C(3) = 1.321(2), C(3)–C(10) = 1.463(2); O(1)–C(1)–C(2) = 120.0(2), O(1)–C(1)–C(4) = 119.9(1), C(2)–C(1)–C(4) = 120.2(1), C(1)–C(2)–C(3) = 121.0(2), C(2)–C(3)–C(10) = 128.4(2). Non-bonded contacts (Å): C(2)⋯C(2') = 5.053(2), C(3)⋯C(3') = 4.162(2) and C(2)⋯C(3') = 4.502(2) (*l*' = 1 – *x*, *y*,  $\frac{1}{2}$  – *z*).



**Fig. 4** CD Spectra of two enantiomeric chiral crystals of compound **1e** in Nujol mulls

Of the four possible dimers of chalcones **1**, three of them, compounds **2**, **4** and **5**, were isolated and their structures were elucidated completely. Since the *syn*-head-to-tail dimer structure of **4** has never been elucidated directly so far, the X-ray



**Fig. 5** Molecular structure of dimer **4a** with crystallographic numbering scheme. Selected bond lengths (Å) and angles (°): O(1)–C(1) = 1.212(2), C(1)–C(2) = 1.512(2), C(1)–C(4) = 1.500(2), C(2)–C(3) = 1.551(2), C(2)–C(3') = 1.586(3), C(3)–C(10) = 1.504(3); O(1)–C(1)–C(2) = 121.5(2), O(1)–C(1)–C(4) = 120.4(2), C(2)–C(1)–C(4) = 118.1(2), C(1)–C(2)–C(3) = 116.1(1), C(1)–C(2)–C(3') = 115.9(1), C(3)–C(2)–C(3') = 91.3(1), C(2)–C(3)–C(2') = 88.7(1), C(2)–C(3)–C(10) = 117.5(1), C(2')–C(3)–C(10) = 118.3(1). The equivalent position ' =  $-x, -y, 2 - z$ .

structure of compound **4a** prepared according to the literature<sup>7</sup> was determined (Fig. 5). However, no *anti*-head-to-tail dimer **7** was detected in any of the photodimerisation experiments which have been carried out so far. This shows that the molecular aggregation such as **8** which gives dimer **7** by photodimerisation would be unfavourable. It is also clarified that mps of chalcone photodimers become higher in the sequence: **4** (232–257 °C) > **2** (164 °C) > **5** (90–139 °C).

## Experimental

Mps were measured on a Yanaco MP-53 apparatus and are uncorrected. <sup>1</sup>H NMR spectra were recorded in CDCl<sub>3</sub> by a JEOL LA-300 300 MHz spectrometer. *J*-Values are given in Hz. IR spectra were recorded on a JASCO FT-IR spectrophotometer for samples as Nujol mulls.

### Photodimerisation of compound **1a** in the molten state

Chalcone **1a** (1 g, 4.8 mmol) was placed in between two Pyrex glass plates and the sample was melted by being heated at 60 °C on a hot-plate, and thereafter was irradiated with a 400 W high-pressure Hg lamp for 24 h. The oily crude product was crystallised by addition of a small amount of MeOH to give, after recrystallisation from MeOH, compound **5a** as prisms (0.31 g, 31%), mp 123–125 °C;  $\nu_{\max}(\text{C}=\text{O})$  1665 cm<sup>-1</sup>;  $\delta_{\text{H}}$  7.8–7.1 (20 H, m, ArH) and 4.61 and 3.97 (2 H, each d, *J* 8.8, CH) (Found: C, 86.39; H, 5.88. C<sub>30</sub>H<sub>24</sub>O<sub>2</sub> requires C, 86.51; H, 5.81%).

### Photodimerisation of compound **1b** in the molten state

By a similar method to that above, chalcone **1b** (1 g, 4.5 mmol) was irradiated in the molten state for 24 h to give, after recrystallisation from MeOH, compound **5b** as prisms (0.27 g, 27%), mp 112–114 °C (lit.,<sup>2</sup> 114 °C);  $\nu_{\max}(\text{C}=\text{O})$  1665 cm<sup>-1</sup>;  $\delta_{\text{H}}$  7.7–7.1 (18 H, m, ArH), 4.57 and 3.95 (2 H, each d, *J* 9.0, CH) and 2.32 (6 H, s, CH<sub>3</sub>).

### Photodimerisation of compound **1c** in the molten state

By a similar method to that above, chalcone **1c** (1 g, 4.1 mmol) was irradiated in the molten state for 24 h to give, after recrystallisation from MeOH, compound **5c** (0.25 g, 25%), mp 128–129 °C;  $\nu_{\max}(\text{C}=\text{O})$  1660 cm<sup>-1</sup>;  $\delta_{\text{H}}$  7.8–7.2 (18 H, m, ArH) and 4.54 and 3.90 (2 H, each d, *J* 9, CH) (Found: C, 74.02; H, 4.33. C<sub>30</sub>H<sub>22</sub>Cl<sub>2</sub>O<sub>2</sub> requires C, 74.23; H, 4.57%).

### Photodimerisation of compound **1d** in the molten state

By a similar method to that above, chalcone **1d** (1 g, 4.2 mmol) was irradiated in the molten state for 24 h to give, after recrystallisation from MeOH, compound **5d** as needles (0.2 g, 20%), mp 90–92 °C;  $\nu_{\max}(\text{C}=\text{O})$  1665 cm<sup>-1</sup>;  $\delta_{\text{H}}$  7.8–7.1 (18 H, m, ArH), 4.58 and 3.88 (2 H, each d, *J* 9.0, CH) and 2.31 (6 H, s, OCH<sub>3</sub>) (Found: C, 80.50; H, 5.63. C<sub>32</sub>H<sub>28</sub>O<sub>4</sub> requires C, 80.65; H, 5.92%).

### Photodimerisation of compound **1c** in the solid state

A stirred suspension of powdered chalcone **1c** (1 g, 4.1 mmol) in water (100 ml) containing a small amount of hexadecyl-(trimethyl)ammonium bromide as a surfactant was irradiated for 12 h at room temp. The reaction mixture was filtered and recrystallised from AcOEt to give compound **4c** as prisms (0.31 g, 31%), mp 255–257 °C;  $\nu_{\max}(\text{C}=\text{O})$  1670 cm<sup>-1</sup>;  $\delta_{\text{H}}$  7.7–7.0 (18 H, m, ArH) and 4.94 and 4.75 (2 H, each dd, *J*<sub>cis</sub> 10.8, *J*<sub>trans</sub> 6.4, CH) (Found: C, 74.13; H, 4.52. C<sub>30</sub>H<sub>22</sub>Cl<sub>2</sub>O<sub>2</sub> requires C, 74.23; H, 4.57%).

### Photodimerisation of compound **1e** in the molten state

By a similar method to that above, chalcone **1e** (1 g, 4.2 mmol) was irradiated in the molten state for 24 h to give, after recrystallisation from MeOH, compound **5e** as prisms (0.32 g, 32%), mp 137–139 °C;  $\nu_{\max}(\text{C}=\text{O})$  1660 cm<sup>-1</sup>;  $\delta_{\text{H}}$  7.7–7.1 (16 H, m, ArH), 4.54 and 3.86 (2 H, each d, *J* 9, CH) and 2.32 and 2.31 (6 H, each s, CH<sub>3</sub>) (Found: C, 86.17; H, 6.39. C<sub>34</sub>H<sub>32</sub>O<sub>2</sub> requires C, 86.41; H, 6.82%).

### X-Ray crystal structure analysis

**Crystal data for compound **4a**.** C<sub>30</sub>H<sub>24</sub>O<sub>2</sub>, *M* = 416.52, triclinic, space group *P*1, *a* = 8.717(3), *b* = 11.863(4), *c* = 6.105(2) Å,  $\alpha$  = 99.67(3)°,  $\beta$  = 105.53(2)°,  $\gamma$  = 107.73(3)°, *V* = 557.5(4) Å<sup>3</sup>, *Z* = 1, *D*<sub>c</sub> = 1.24 g cm<sup>-3</sup>, crystal dimensions 0.38 × 0.38 × 0.20 mm,  $\mu(\text{Mo-K}\alpha)$  = 0.76 cm<sup>-1</sup>. Diffraction data were collected on a Rigaku AFC-7R diffractometer with graphite-monochromated Mo-K $\alpha$  radiation ( $\lambda$  = 0.710 73 Å) to  $2\theta_{\max}$  = 55°. Intensity data were corrected for Lorentz and polarisation effects. All calculations were carried out using the TEXSAN crystallographic software package from Molecular Structure Corporation. Non-hydrogen and hydrogen atoms were refined anisotropically and isotropically, respectively. The final cycle of full-matrix least-squares refinement was based on 1530 observed reflections [*I* > 3 $\sigma$ (*I*)] and 194 parameters and converged with *R* = 0.046, *R*<sub>w</sub> = 0.060 (*w* = [ $\sigma^2(F_o) + 0.0009F_o^2$ ]), and GOF = 1.44;  $\Delta\rho_{\max}$  = 0.23 e Å<sup>-3</sup>.

**Crystal data for compound **5a**.** C<sub>30</sub>H<sub>24</sub>O<sub>2</sub>, *M* = 416.52, monoclinic, space group *P*2<sub>1</sub>/*c*, *a* = 10.905(2), *b* = 9.672(2), *c* = 21.406(2) Å,  $\beta$  = 90.79(1)°, *V* = 2257.5(5) Å<sup>3</sup>, *Z* = 4, *D*<sub>c</sub> = 1.225 g cm<sup>-3</sup>, crystal dimensions 0.32 × 0.32 × 0.20 mm,  $\mu(\text{Mo-K}\alpha)$  = 0.75 cm<sup>-1</sup>. Data collection and structure analysis were performed by the same method as described for compound **4a**. *R* = 0.039, *R*<sub>w</sub> = 0.058, GOF = 1.31 for 2465 observed reflections [*I* > 3 $\sigma$ (*I*)] and 386 parameters;  $\Delta\rho_{\max}$  = 0.14 e Å<sup>-3</sup>.

**Crystal data for compound **1b**.** C<sub>16</sub>H<sub>14</sub>O, *M* = 222.29, monoclinic, space group *C*2/*c*, *a* = 14.976(4), *b* = 9.843(3), *c* = 17.561(3) Å,  $\beta$  = 105.83(2)°, *V* = 2490(1) Å<sup>3</sup>, *Z* = 8, *D*<sub>c</sub> = 1.186 g cm<sup>-3</sup>, crystal dimensions 0.60 × 0.50 × 0.53 mm,  $\mu(\text{Mo-K}\alpha)$  = 0.72 cm<sup>-1</sup>. Data collection and structure analysis were performed by the same method as described for compound **4a**. *R* = 0.042, *R*<sub>w</sub> = 0.054, GOF = 2.83 for 1915 observed reflections [*I* > 3 $\sigma$ (*I*)] and 155 parameters;  $\Delta\rho_{\max}$  = 0.15 e Å<sup>-3</sup>.

**Crystal data for compound **1c**.** C<sub>15</sub>H<sub>11</sub>ClO, *M* = 242.70, triclinic, space group *P*1, *a* = 7.565(2), *b* = 14.060(2), *c* = 5.858(1) Å,  $\alpha$  = 90.09(2)°,  $\beta$  = 92.15(2)°,  $\gamma$  = 81.70(2)°, *V* = 616.0(3) Å<sup>3</sup>, *Z* = 2, *D*<sub>c</sub> = 1.308 g cm<sup>-3</sup>, crystal dimensions 0.30 × 0.14 × 0.14 mm,  $\mu(\text{Mo-K}\alpha)$  = 2.88 cm<sup>-1</sup>. Data collection and structure analysis were performed by the same method as described for compound **4a**. *R* = 0.034, *R*<sub>w</sub> = 0.049, GOF = 1.11 for 1241

observed reflections [ $I > 3\sigma(I)$ ] and 199 parameters;  $\Delta\rho_{\max} = 0.13 \text{ e \AA}^{-3}$ .

Full crystallographic details, excluding structure factor tables, have been deposited at the Cambridge Crystallographic Data Centre (CCDC). For details of the deposition scheme, see 'Instructions for Authors', *J. Chem. Soc., Perkin Trans. 1*, available via the RSC Web page (<http://www.rsc.org/authors>). Any request to the CCDC for this material should quote the full literature citation and the reference number 207/186.

### References

- 1 *Org. Synth.*, 1941, **1**, 78; F. Toda, K. Tanaka and K. Hamai, *J. Chem. Soc., Perkin Trans. 1*, 1990, 3207.
- 2 H. Stobbe and K. Brenner, *J. Prakt. Chem.*, 1929, **123**, 241.
- 3 D. Rabinovich and G. M. J. Schmidt, *J. Chem. Soc. B*, 1970, 6.
- 4 G. M. J. Schmidt, Photochemistry of the excited state, in *Reactivity of the Photoexcited Organic Molecules*, J. Wiley, New York, 1967.
- 5 H. Stobbe and A. Heusel, *Ber. Dtsch. Chem. Ges.*, 1926, **59**, 2254.
- 6 G. M. J. Schmidt, *Pure Appl. Chem.*, 1971, **27**, 647.
- 7 K. Tanaka and F. Toda, *J. Chem. Soc., Chem. Commun.*, 1983, 593.
- 8 M. Kaftory, K. Tanaka and F. Toda, *J. Org. Chem.*, 1985, **50**, 2154.
- 9 K. Ohkuma, S. Kashino and M. Haida, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 627.
- 10 D. Rabinovich, *J. Chem. Soc. B*, 1970, 11.
- 11 S. Caccamese, J. A. McMillan and G. Montaudo, *J. Org. Chem.*, 1978, **43**, 2703.
- 12 C. F. H. Allen and G. F. Frame, *Can. J. Res.*, 1932, **6**, 605.
- 13 D. Rabinovich and Z. Shakked, *Acta Crystallogr., Sect. B*, 1974, **30**, 2829.
- 14 K. Penzien and G. M. Schmidt, *Angew. Chem., Int. Ed. Engl.*, 1969, **8**, 608.
- 15 F. Toda, H. Miyamoto, S. Kikuchi, R. Kuroda and F. Nagami, *J. Am. Chem. Soc.*, 1996, **118**, 11 315; F. Toda, H. Miyamoto and K. Kanemoto, *J. Org. Chem.*, 1996, **61**, 6490; H. Koshima, T. Nakagawa, T. Matsuura, H. Miyamoto and F. Toda, *J. Org. Chem.*, 1997, **62**, 6322.

Paper 7/07380A

Received 13th October 1997

Accepted 12th January 1998